

Theory of the Resonant Neutron Scattering of High T_c Superconductors

Eugene Demler and Shou-Cheng Zhang

Department of Physics, Stanford University, Stanford, CA 94305

(December 20, 1994)

Recent polarized neutron scattering experiments on $YBa_2Cu_3O_7$ have revealed a sharp spectral peak at the (π, π) in reciprocal lattice centered around the energy transfer of 41 meV. We offer a theoretical explanation of this remarkable experiment in terms of a new collective mode in the particle particle channel of the Hubbard model. This collective mode yields valuable information about the symmetry of the superconducting gap.

PACS numbers: 74.72.Bk, 61.12.Bt, 61.12.Ex

Recently, both unpolarized and polarized neutron scattering experiments have been performed on the $YBa_2Cu_3O_7$ high T_c superconductors [1–3]. In particular, polarized neutron experiment [2] shows an extremely sharp spectral feature in the spin flip channel. This feature is centered around (π, π) in reciprocal space, and peaked at 41 meV with a width narrower than the instrumental resolution. This feature also has an interesting temperature dependence. In the experiment by Mook *et al*, while it exists above the superconducting transition temperature of $T_c = 92.4K$, its intensity scales like the superfluid density below the transition. More recently, Fong *et al* performed detailed spin unpolarized neutron experiments with a careful subtraction of the phonon background. They found that the 41 meV mode disappears above the superconducting transition temperature.

In this letter, we offer a theoretical explanation of this remarkable experiment. We first show that for a general class of tight binding Hamiltonian including the Hubbard and the $t - J$ model, there exist well-defined collective modes in the particle particle channel centered around momentum (π, π) . The spin quantum number of this excitation can either be a singlet or a triplet. The singlet excitation has been discussed by Yang [4] and one of us [5], and is in fact an exact eigenstate of the Hubbard model. Normally, collective excitations in the particle particle channel are inaccessible experimentally. However, one of us [5] argued that if the ground state of the model in consideration is superconducting, one can couple to it through a particle hole excitation, because the BCS condensate is a coherent mixture of particles and holes. Based on this consideration, one of us [5] predicted a possible new collective mode of the high T_c superconductors. It is a spin singlet excitation peaked at (π, π) , has a well defined energy of $U - 2\mu$ and its intensity scales like the superfluid density. Possibly because it is hard to distinguish it from other excitations in the system, this mode has not yet been detected experimentally.

However, the basic arguments can be easily generalized from the singlet to the triplet case. Besides the above mentioned collective mode in the singlet particle particle

channel, there also exists a well defined collective mode in the triplet channel near total momentum (π, π) . This is true for a large class of tight binding models, like the Hubbard or the $t - J$ model. The energy spectrum of a non-interacting pair of particles or holes generally consists of a continuum labeled by their relative momentum. However, for tight binding models, this continuum collapses to a point when the total momentum of the pair is (π, π) . (See Figure 1).




FIG. 1. The energy spectrum along the (π, π) direction. The dots correspond to the antibonding state and the solid line is the edge of the continuous spectrum. Here the numerical calculations were done for $J = \tilde{t}$ and $n = 0.85$. As δ we denoted $\frac{\pi}{50}$.

A triplet pair generally has a repulsive interaction when placed on the neighboring site. Because of the collapse of the particle particle continuum, this repulsive interaction leads to an *anti-bound* state near total momentum $(\pi, \pi, \dots)$ in any space dimensions. (See Figure 1). This anti-bound triplet state manifests itself as a collective excitation of the many-body system, or as a pole in the Greens function of the particle particle channel. Most physical probes do not couple to this channel. However, if the model in consideration is superconducting (which is an assumption of our theory), then a spin flip scattering of the neutron can couple directly to the

triplet particle particle excitation. Physically this process is nothing but the spin flip scattering of a Cooper pair in the RCS condensate. There is an overwhelming evidence that the Cooper pair in the high T_c materials is a spin singlet with total momentum zero [6]. Therefore a spin flip scattering of the Cooper pair creates a triplet state. The orbital angular momentum or the parity selection rule forbids such a transition if the momentum loss of the neutron is zero, on the other hand, it can be shown simply that the matrix element of this coupling is maximal when the momentum loss of the neutron is at (π, π) .

Based on the above reasoning, we shall interpret the sharp spectral feature observed in the polarized neutron experiment in terms of the triplet collective mode in the particle particle channel. We begin with some analytic calculations of the properties of this collective mode and its contribution to the dynamical spin spin correlation function. Subsequently, we compare our predictions with a number of characteristic features of the resonance observed in experiments. Finally we point out some difficulties with our interpretation and their possible resolution.

Our anti-bound state is different from the excitons of a superconductor considered by Bardasis and Schrieffer [7]. These excitons form because of an *attractive* potential in a given angular momentum channel, and they exist inside the superconducting gap and near total momentum $q = 0$. Our model is also different from the magnetic susceptibility in the superconducting state computed using RPA [8], since it involves multiple scattering in the particle hole channel. More recently, Bulut and Scalapino [9] considered a model of the bi-layer superconductors and argued that a dynamic nesting from the bonding to anti-bonding Fermi surface could give rise to a collective resonance at (π, π) . Our main difference lies in the fact that their resonance is in the particle hole channel and as such it can exist well above T_c . The bi-layer band structure plays a crucial role in their model while it is irrelevant in our case.

We consider the following model defined on a two dimensional square lattice:

$$\mathcal{H} = -t \sum_{\langle ij \rangle} c_{i\sigma}^\dagger c_{j\sigma} + J \sum_{\langle ij \rangle} \vec{S}_i \cdot \vec{S}_j + U \sum_i n_{i\uparrow} n_{i\downarrow} - \mu \sum_i c_{i\sigma}^\dagger c_{i\sigma} \quad (1)$$

In the limit $U \rightarrow \infty$, we recover the $t - J$ model. On the other hand, if we keep U small, we can regard the above model as an effective Hamiltonian of the weak coupling Hubbard model, where the J term arises from the paramagnon interaction on the nearest neighbor sites. To study the spectrum in the triplet particle particle channel, we consider the operator

$$O_q^\dagger = \sum_p f_q(p) c_{p+q\uparrow}^\dagger c_{-p\uparrow}^\dagger \quad (2)$$

which describes a pair of particles with center of mass momentum q and relative momentum p . The equation of

motion for this operator is given by its commutator with the Hamiltonian, $[\mathcal{H}, O_q^\dagger]$. If we factorize the resulting commutator in terms of $O_q^\dagger$ and the expectation values of the density $n_{k\sigma}$, we obtain:

$$\begin{aligned} [\mathcal{H}, O_q^\dagger] = & \sum_p (\epsilon_{p+q} + \epsilon_{-p}) f_q(p) c_{p+q\uparrow}^\dagger c_{-p\uparrow}^\dagger + 2(U n_{\downarrow} - \mu) O_q^\dagger \\ & + \frac{J}{8} \sum_p c_{p+q\uparrow}^\dagger c_{-p\uparrow}^\dagger \times \frac{1}{N} \sum_{p'} f_q(p') \\ & \times (\eta(p - p') - \eta(p + p' + q)) (1 - n_{p'-q\uparrow} - n_{p'\uparrow}) \\ & - \frac{3J}{8} \sum_p f_q(p) c_{p+q\uparrow}^\dagger c_{-p\uparrow}^\dagger \\ & \times \frac{1}{N} \sum_{p'} n_{p'} (\eta(p + p') + \eta(p' - p - q)) \quad (3) \end{aligned}$$

where $n_\sigma = \frac{1}{N} \sum_k n_{k\sigma}$ is the average density of electrons with spin σ , N is the total number of lattice sites and $\eta(p) = \sum_{\vec{a}} \exp[i\vec{p} \cdot \vec{a}]$ is a geometrical factor coming from the summation over the nearest neighbors. In this equation, the first term describes the kinetic energy of the pair of particles in consideration. For tight binding models with nearest neighbor hopping, $\epsilon_k = -2t(\cos k_x + \cos k_y)$. In this case, the kinetic energy of the pair vanishes when the total momentum $q = (\pi, \pi)$. The second term describes the Hartree interaction of the spin up pair with the average density of the down spins in the background and the chemical potential energy of adding a pair of particles. In the large U limit, the leading contribution to the chemical potential is given by $Un/2$. In fact, for Hubbard model at half filling, *i.e.* at $n = 1$, $\mu = U/2$ is an exact relation. Therefore, the second terms cancels in the leading order in U , and reaches a finite limit as $U \rightarrow \infty$. The third term gives the multiple scattering of the particles with each other on a restricted phase space due to the filled Fermi sea. The last term describes the Fock self-energy of the quasi-particles. The method of factorizing the operator equation of motion is equivalent to the T matrix approximation in the diagrammatic calculations, it is exact in the low density limit. The collective mode for the triplet particle particle excitation is obtained by equating the right hand side of (3) to $E_q O_q^\dagger$, the resulting eigenvalue equation is given by

$$1 = \frac{J}{2N} \sum_p \frac{\sin^2(p_\alpha) (1 - n_{\frac{q}{2}-p\uparrow} - n_{\frac{q}{2}+p\uparrow})}{\tilde{E}_q^\alpha + 4\tilde{t} [\cos(p_x) \cos(\frac{q_x}{2}) + \cos(p_y) \cos(\frac{q_y}{2})]} \quad (4)$$

with α being x or y , $\tilde{t} = t + \frac{3J}{4N} \sum_p n_{p\downarrow} \cos(p_x)$ and $E_q = \tilde{E}_q + 2(U n_{\downarrow} - \mu)$. It is straightforward to see that a discrete eigenvalue exists at $q = (\pi, \pi)$, with energy of

$$\tilde{E} = \frac{J}{4} (1 - \frac{2}{N} \sum_p n_p \cos^2 p_x) \quad (5)$$

Discrete eigenvalues also exist for a finite range in q space. There are in general two eigenvalues for each q , one corresponds to a p wave pair oriented in the x direction,

while the other is oriented in the y direction. The two eigenvalues are degenerate only along the diagonal when $q_x = q_y$. The above eigenvalue equation can easily be solved numerically and the dispersion of the collective mode is shown in figure 1. $\tilde{t}$ is basically the renormalized hopping matrix element of the quasi-particle. The range in momentum space over which the collective mode exists depends on the ratio of $J/\tilde{t}$. Numerous numerical calculations indicate that the quasi-particle bandwidth is of the order of J [10]. Here we take a semi-phenomenological approach and choose $\tilde{t}$ rather than t as a free parameter. For a ratio of $J/\tilde{t} = 1$, and $n = 0.85$ we see that the collective mode exists over $\frac{\pi}{50}$ of the momentum space.

In the experiments with the polarized neutrons scattering one measures the dynamic spin-spin correlation function

$$S(q, \omega) = \sum_n |\langle n | S_q^\dagger | 0 \rangle|^2 \delta(\omega - \omega_{n0}) \quad (6)$$

where $|0\rangle$ and $|n\rangle$ are the ground and excited states of the system and $S_q^\dagger = \sum_p c_{p+q\uparrow}^\dagger c_{p\downarrow}$. Using the operator equation $[\mathcal{H}, O_{q,\alpha}^\dagger] = E_{q,\alpha}^\alpha O_{q,\alpha}^\dagger$ we can construct a class of approximate excited states of the Hubbard Hamiltonian as $|n\rangle = O_{q,\alpha}^\dagger |0\rangle$. The same operator equation for $O_{q,\alpha}$ shows that $O_{q,\alpha} |0\rangle = 0$. With these relations, we can calculate the contribution of this approximate eigenstate to $S(q, \omega)$ at zero temperature:

$$\begin{aligned} S_0(q, \omega) &= \\ &= \sum_\alpha \langle 0 | O_{q,\alpha} S_q^\dagger | 0 \rangle^2 \delta(\omega - E_q^\alpha) + \sum_{n'} \\ &= \sum_\alpha \langle 0 | [O_{q,\alpha}, S_q^\dagger] | 0 \rangle^2 \delta(\omega - E_q^\alpha) + \sum_{n'} \end{aligned} \quad (7)$$

where $\sum_{n'}$ denotes the contribution from states other than $O_{q,\alpha}^\dagger |0\rangle$, and we used the fact that $O_{q,\alpha} |0\rangle = 0$ to replace the product of two operators by their commutator. Evaluating the commutator we obtain

$$\begin{aligned} S_0(q, \omega) &= \\ &= \sum_\alpha \langle 0 | 2 \sum_p f_q^{(\alpha)} (p - \frac{q}{2}) c_{-p+\frac{q}{2}\uparrow} c_{p-\frac{q}{2}\downarrow} | 0 \rangle^2 \delta(\omega - E_q^\alpha) \\ &= 4 \sum_\alpha |\sum_p f_q^{(\alpha)} (p - \frac{q}{2}) \Delta_{p-q/2}|^2 \delta(\omega - E_q^\alpha) \end{aligned} \quad (8)$$

Summation over α corresponds to the two kinds of antibonding solutions. The overlap matrix element is finite for both extended s wave and d wave pairing in the ground state. If we take the d wave symmetry of the order parameter $\Delta_k = \Delta_0(\cos(k_x) - \cos(k_y))$ and, for example, $f_q^{(x)}$ - a p wave oriented in the x direction. Then,

$$\begin{aligned} &\sum_p f_q^{(x)} (p - \frac{q}{2}) \Delta_{p-q/2} = \\ &= \frac{1}{A_q} \sum_p \frac{\sin(p_x) \Delta_{p-q/2} (1 - n_{\frac{q}{2}-p} - n_{\frac{q}{2}+p})}{\tilde{E}_q + 4\tilde{t}(\cos(p_x) \cos(\frac{q_x}{2}) + \cos(p_y) \cos(\frac{q_y}{2}))} = \\ &= -\frac{2\Delta_0}{JA_q} \sin(\frac{q_x}{2}) \end{aligned} \quad (9)$$

where A_q is the normalization factor of the wave function

$$A_q^2 = \sum_p \frac{\sin^2(p_x) (1 - n_{\frac{q}{2}-p} - n_{\frac{q}{2}+p})^2}{[\tilde{E}_q + 4\tilde{t}(\cos(p_x) \cos(\frac{q_x}{2}) + \cos(p_y) \cos(\frac{q_y}{2}))]^2}$$

Finally we have

$$\begin{aligned} S(q, \omega, T) &= 16 \frac{\Delta_0^2(T)}{J^2 A_q^2} [\sin^2(\frac{q_x}{2}) \delta(\omega - E_q^x) \\ &+ \sin^2(\frac{q_y}{2}) \delta(\omega - E_q^y)] \left(1 + n_B(\frac{\omega}{T})\right) \end{aligned} \quad (10)$$

where the last factor takes care of the finite temperature in spin-spin correlation functions. One can easily see though, that the main temperature dependence of $S(q, \omega, T)$ comes from $\Delta_0(T)$. We also see that the intensity vanishes at zero external momentum and is maximal at (π, π) . This fact can simply be attributed to the conservation of orbital angular momentum. Since the Cooper pair in the ground state has angular momentum either $l = 0$ or $l = 2$, a finite momentum transfer is required for it to be excited to a state with angular momentum $l = 1$. The dispersion of two states is not very big giving rise to only a small splitting of two levels which never exceeds 15%.

We also see that the spectral intensity is simply proportional to the BCS order parameter or the superfluid density. This is so because the BCS order parameter provides the coupling from the particle hole channel to the particle particle channel [5]. This extra spectral weight at energy $E_{q,\alpha}$ is transferred from the low energy sector, since the singlet BCS pairing removes the low energy spin fluctuation. Our theory is consistent with the fact that the 41meV peak intensity seems to scale with the superfluid density below T_c . This conclusion agrees with the experimental results obtained by Fong *et al* [3], but it is inconsistent with the fact that this mode does not vanish above T_c in the experiment by Mook *et al* [2]. At this moment, it is an experimental question that has to be resolved. We also note that our calculation is not fully self-consistent, and assumes that the mode energy does not change significantly when the system becomes superconducting. This is true for a variety of collective modes like the plasmon and the singlet particle particle mode [4,5] (the so called η pair). But we do not have a general proof in the present case. A fully self-consistent calculation will be carried out in the future [11], but we anticipate that the basic physics remain unchanged.

All the above discussions were restricted to the two dimensional CuO plane, which we model by the two dimensional Hubbard model. However, it can be simply generalized to three dimensions. If one takes a three dimensional Hubbard model, one finds that collective mode exists near (π, π, π) rather than $(\pi, \pi, 0)$. This is consistent with the experiment where the third component of the momentum transfer is also π .

The fact that the particle particle collective mode always exists at $(\pi, \pi, \dots)$ is a special property of the tight

binding model on a bi-partite lattice. How would a next-nearest neighbor hopping term change the results? In this case, the dispersion relation is given by

$$\epsilon_k = -2\tilde{t}(\cos k_x + \cos k_y) - 2t'(\cos(k_x + k_y) + \cos(k_x - k_y))$$

where t' denotes the amplitude of the next-nearest neighbor hopping. The x and the y collective modes are mixed into the bonding $\sin(p_x) + \sin(p_y)$ and the antibonding $\sin(p_x) - \sin(p_y)$ combination, and we obtain a matrix equation for the eigenvalues of these collective modes:

$$\det \left\| \frac{J}{2} \frac{1}{N} \sum_p \frac{\sin(p_\alpha) \sin(p_\beta) z_{pq}}{\tilde{E} - \Omega_{pq}} - \delta_{\alpha\beta} \right\| = 0 \quad (11)$$

where $z_{pq} = 1 - n_{\frac{q}{2}-p} - n_{\frac{q}{2}+p}$ and $\Omega_{pq} = \epsilon_{\frac{q}{2}-p} + \epsilon_{\frac{q}{2}+p}$. It is easy to see that the antibonding state $E_{\pi,\pi}^-$ always exists, while there is a critical coupling

$$t'_{cr} = 1.2 \times 10^{-4} J. \quad (12)$$

for the bonding state, so that for $t' > t'_{cr}$, it ceases to exist at (π, π) . In real experiments t' and J are of the same order of magnitude, therefore, one can safely conclude that the anti-bonding state disappears into the continuum. This conclusion leads directly to a measurement of the *symmetry of the superconducting gap*. Below the superconducting transition temperature, the intensity of the anti-bonding state as measured in the neutron scattering experiment is proportional to

$$\left| \sum_p f_q^{(\alpha)}(p - \frac{q}{2}) \Delta_{p-q/2} \right|^2 \quad (13)$$

where $f_q^{(\alpha)}(p - \frac{q}{2}) \propto \sin(p_x) - \sin(p_y)$. From this equation we see immediately that the intensity is non vanishing if and only if the gap symmetry is of the d wave type. We therefore argue that the existence of the neutron resonance determines the *symmetry* of the pairing gap of the high T_c superconductors to be of the d wave type, consistent with the theories where pairing interaction arises from the spin fluctuations [12–14]. It could also be consistent with more exotic possibilities of $d_{x^2-y^2} + id_{xy}$ pairing symmetry [15].

We conclude that the basic features of the observed polarized neutron scattering experiment can be explained in terms of a new particle particle collective mode in the Hubbard model. The energy of the mode at (π, π) is given by (5) in the case of $t' = 0$ and implicitly by (11) for finite t' . Unfortunately, the energy of the mode depends on all three parameters μ , J and t' . While these parameters can be fitted to the experimental value, the fitting itself is not a confirmation of our theory. The mode is centered around (π, π, π) for two different reasons, both because the particle particle continuum at this momentum is minimal so that the anti-bound state could exist, and because of the conservation of angular momentum for exciting a singlet Cooper pair to a triplet state. This is exactly the momentum transfer of the excitation observed in experiment. The intensity of the mode scales

with the superfluid density because the BCS pairing amplitude is involved in converting a particle hole pair into a particle particle pair. Above T_c , our simple calculation shows that the mode disappears, a much more sophisticated calculation is required to understand if the fluctuation pairing effects could give rise to a resonance even above T_c . When the effects of next-nearest-neighbor hopping is included, only the anti-bonding $\sin(p_x) - \sin(p_y)$ mode can exist and the region in momentum space for the existence of this collective mode becomes much narrower. However, the sharp collective mode might still exist as a broad resonance outside of this region. The antibonding collective mode only has an overlap with the d wave order parameter, and we conclude that the experimental observation of the collective resonance in the neutron scattering experiment can only be consistent with the d wave symmetry of the pairing gap.

We are extremely grateful to Prof. D. Scalapino for a stimulating journal club talk on the experiment and generous sharing of his ideas on the problem. We would also like to thank Prof. R. B. Laughlin, V. Emery, N. Bulut and Z. X. Shen for many enlightening discussion on the experiment. Part of this work is supported by the Center for Materials Research at Stanford University.

-
- [1] J. Rossat-Mignod, *et al*, *Physica* **180B**, 383 (1992).
 - [2] H. A. Mook, *et al*, *Phys. Rev. Lett.* **70**, 3490 (1993).
 - [3] H. F. Fong *et al*, Princeton University preprint.
 - [4] C. N. Yang, *Phys. Rev. Lett.* **63**, 2144 (1989); C.N. Yang and S.C. Zhang, *Mod. Phys. Lett.* **B4**, 759 (1990).
 - [5] S. C. Zhang, *Phys. Rev. Lett.* **65**, 120 (1990); S. C. Zhang, *Int. J. Mod. Phys.* **B5**, 153 (1991).
 - [6] D. J. Scalapino, *Phys. Rep.* to be published. *Phys. Rev.* **121**, 1050, (1961).
 - [7] A. Bardasis and J. R. Schrieffer, *Phys. Rev.* **121**, 1050, (1961).
 - [8] P. Monthoux and D. J. Scalapino, *Phys. Rev. Lett.* **72**, 1874 (1994).
 - [9] N. Bulut and D. J. Scalapino, UCSB preprint.
 - [10] E. Dagotto, *Rev. Mod. Phys.* **66**, 763 (1994).
 - [11] N. Bulut, E. Demler, D. J. Scalapino and S. C. Zhang, work in progress.
 - [12] D. J. Scalapino, J. E. Hirsch and E. Y. Loh, *Phys. Rev.* **B 34**, 8190 (1986).
 - [13] J. R. Schrieffer, X. G. Wen and S. C. Zhang, *Phys. Rev.* **B 41**, 6399 (1990).
 - [14] P. Monthoux, A. V. Balatsky and D. Pines, *Phys. Rev.* **B46**, 14803 (1992).
 - [15] R. B. Laughlin, Stanford University preprint.